

Razowski, J. 1983. Motyle (Lepidoptera) Polski. Cz. VI—Olethreutinae: Olethreutidii. *Monogr. Fauny Polski* 13: 1–177 [In Polish].

A tale of two galls.—Galls are distinctive and abnormal growths produced by a plant in response to the influence of an organism (Redfern & Askew, 1992). Frequently, galls are lush outgrowths and have the appearance of being potentially attractive food for other species. I have previously shown that some free-living psyllids and aphids can have faster growth rates when feeding on buckthorn (*Rhamnus catharticus* L.) leaves with galls formed by *Trichohermes walkeri* Förster, and that these different Homoptera show associations ranging from obligate use of galled leaves, through showing a preference for feeding on galled leaves, to showing no significant association with *T. walkeri* galls (McLean, 1994).

Although such feeding on galled leaves by sap-sucking Homoptera may have some negative effects on a gall-former such as *T. walkeri* (though this has yet to be tested for this species), eating gall tissue, and/or eating the gall-inducing organism within the gall, is obviously more directly damaging to the gall-former. I have recently observed two instances of interactions between galls and free-living species which offer contrasting outcomes for the respective gall-formers.

First, I have seen a grey squirrel, *Sciurus carolinensis* Gmelin, feeding on the contents of leaf-petiole galls formed by the aphid *Pemphigus spirothecae* Passerini (Homoptera: Pemphigidae) on Lombardy poplar *Populus nigra* L. var. *italica* at the rear of Monkstone House, City Road, Peterborough at around 19.30 BST on 15 August and again at about 18.30 BST on 20 August 1996. The debris of broken galls and discarded leaves rained down at the rate of several per minute, but whether it was the honeydew, the insects themselves, or these items together which were the principal attraction for the squirrel (or squirrels, as I could not tell whether the same individual was responsible on both dates) remains a mystery. However, examination of the opened galls suggested that it was the gall contents rather than the galls themselves which were consumed by the squirrel(s), which had perhaps acquired a sweet tooth!

Second, in my garden at Miller Way there is a female willow bush, *Salix aurita* L. on which I noticed an inflorescence gall, possibly formed by the mite *Phytopus triradiatus* (Nalepa) (Acari: Eriophyidae) according to the figures in Redfern & Askew (1992) and Stubbs (1986), though the overall form of the gall was more compact. On 20 August 1996 I saw that a substantial number of higher branches on the bush had been defoliated by the clutch-feeding larvae of the buff-tip, *Phalera bucephala* L. (Lepidoptera: Notodontidae). The branch with the single inflorescence gall was completely defoliated around the gall, but the gall itself was intact. Whether these voracious larvae ignored the gall because of its non-leaf shape and texture, and/or whether some chemical(s) deterred feeding is unknown. Clearly, there would be a strong selective advantage favouring those galls which are distasteful to chewing insects such as Lepidoptera larvae (not to mention other browsers, including mammals) but whether many galls are unpalatable seems unclear, and is worth further observation and experiment.—IAN F. G. MCLEAN, 109 Miller Way, Brampton, Huntingdon, Cambs PE18 8TZ.

REFERENCES

- McLean, I. F. G. 1994. Interactions between *Trichohermes walkeri* (Homoptera: Psylloidea) and other Homoptera on *Rhamnus catharticus*. In: Williams, M. A. J. (Ed.), *Plant galls*. Systematics Association Special Volume No. 49, Clarendon Press, Oxford, pp. 151–160.

Redfern, M. & Askew, R. R. 1992. *Plant galls*. Naturalists' Handbooks 17, Richmond Publishing Co., Slough.

Stubbs, F. (Ed.) 1986. *Provisional keys to British plant galls*. British Plant Gall Society, Yorkshire.

***Ampedus sanguinolentus* (Schrank) (Coleoptera: Elateridae) on Wimbledon Common and Putney Heath.**—Mr Maxwell Barclay found this species at Wimbledon Common in October 1995. On 14.iv.1996 I found a specimen of the same species on Putney Heath, part of the same area of heathland, under the bark of a fallen birch, a little south of King's Mere.

Mr A. A. Allen and Mr B. A. Cooper have previously recorded this species on Wimbledon Common, in March 1946, and 1966 (Hodge, 1993), but I know of no other records for it since then in this locality. *Ampedus sanguinolentus* is graded a "notable A" species, and is restricted to heathland in southern England. It has an association with rotten birch but may develop in other species, for example pine and sallow.

I wish to thank Mr Maxwell Barclay, who alerted me to the presence of this species on the Common—MARTIN HENDERSON, 18 Landseer Close, Merton Abbey, London SW19 2UT.

REFERENCE

Hodge, P. 1993. *An invertebrate survey of the grassland, heathland and wetland habitats of the eastern half of Wimbledon Common*. Unpublished report for English Nature, Peterborough.

ANNOUNCEMENT

BENHS on the Internet—The Society now has an official World Wide Web page on the Internet. The site is linked to and from other entomological and natural history sites in the United Kingdom and the United States.

Only basic information on the Society, mainly a reproduction of information given in advertising material and sales literature and already made available to members, has been included at this stage. This information includes:

- an introduction to the Society and its aims and objectives,
- details of indoor meetings and the annual exhibition (but not field meetings and workshops),
- Society publications for sale.

As there is no control over who accesses this information, Council have decided that sensitive information concerning the Society's assets and some activities should not be made available on the Internet. Members who might wish to mention the Society on other sites are asked to approach Council before publishing information by this means.

The Society's URL is: <http://ourworld.compuserve.com/homepages/pyo/BENHS.htm>.